

Medicine <i>Trade name</i>	Strength <i>(powder volume)</i> [volume]	IV administration	Other routes of administration	Compatible IV fluids	Additional information
Papaverine <i>DBL Papaverine</i>	12 mg/mL [10 mL] 30 mg/mL [1 mL]	Give undiluted over at least 1 – 2 minutes.	IM: Suitable. SC: Not recommended.	Sodium chloride 0.9% Y-site only and papaverine 5 mg/mL or weaker: Glucose 5% Glucose 5% and sodium chloride 0.9% Potassium chloride 20 mmol/L	Caution: rapid IV administration may cause arrhythmias or apnoea. Protect from light during storage.
Paracetamol <i>Paracetamol (Actavis)</i> <i>Paracetamol (AFT)</i> <i>Paracetamol (Pfizer)</i> <i>Paracetamol (Kabi)</i> <i>Perfalgan</i>	10 mg/mL [50 mL, 100 mL]	Use undiluted or dilute to 1 mg/mL and infuse over 15 minutes.	IM: Not recommended. SC: Not recommended.	Glucose 5% Glucose 5% and sodium chloride 0.9% Glucose 10% Hartmann's Sodium chloride 0.9% Y-site only: Plasma-Lyte 148 Potassium chloride 20 mmol/L	Use diluted solution within one hour of preparation (including infusion time). Discard any unused vial contents. Protect from light during storage.